

Recurrence of atrial fibrillation after switching from brand to generic atenolol

Luca Gallelli, Francesca Maida, Orietta Staltari, Pierandrea Rende, Emilio Russo, Benedetto Caroleo, Giovambattista De Sarro

Department of Health Science, University of Catanzaro, Catanzaro, Italy

Received: 23-01-2014

Revised: 31-05-2014

Accepted: 28-06-2014

ABSTRACT

Beta blockers are the initial treatment for rate control of supraventricular tachyarrhythmia in patients without a history of myocardial infarction or left ventricular dysfunction. In this article we report the recurrence of atrial fibrillation after switching to the generic formulation of atenolol.

Key words: Atenolol, atrial fibrillation, brand drug, clinical ineffectiveness, generic drug

INTRODUCTION

Supraventricular arrhythmias are common. They are more common in women than in men.^[1,2]

Often β -adrenoceptor antagonists (β -blockers) are able to control the ventricular rate by slowing the atrioventricular conduction.^[3]

Several years ago, generic drugs were introduced in Italy in agreement with the Finance Law of 1996 (Law n. 549/1995 in G.U. n. 302 of 29.12.1995). These formulations are equivalent to brand one if they have the same active substance (with a difference of $\pm 5\%$), the same pharmaceutical form, the same therapeutic indications, and a similar bioequivalence ($\pm 20\%$) with the reference medicinal product (Law n. 425/1996 in

G.U. n. 208 of 05.09.1996; Legislative Decree No. 219/06). In this article we report recurrence of atrial fibrillation after changing from the brand to the generic formulation of atenolol.

CASE REPORT

A 63-year-old woman, with a past medical history of paroxysmal atrial fibrillation, was in sinus rhythm, on maintenance beta blocker (Tenormin) therapy. She presented with symptomatic palpitations, nausea, weakness, and dizziness. Clinical evaluation revealed the presence of tachycardia (heart rate of 130 heart beats and atrial fibrillation by electrocardiography). Her blood pressure was 130/80 mmHg, without clinical or laboratory features of heart failure.

History revealed that about 18 years ago (age of 45 years), she was treated for hypertension and hyperthyroidism and at the time of this presentation the patient was treated with telmisartan (Micardis[®], 40 mg/day), amlodipine (Norvasc[®], 5 mg/day), and tapazole (Tiamazole[®], 5 mg/day). Moreover, in 2002, she started atenolol (Tenormin[®] 50 mg bid) for the development of tachycardia, with a good control of heart symptoms. In-depth evaluation revealed that in March 2013, the pharmacist switched the medication from the brand

Access this article online	
Quick Response Code:	Website: www.jpharmacol.com
	DOI: 10.4103/0976-500X.149146

Address for correspondence:

Luca Gallelli, Department of Health Science, School of Medicine, University of Catanzaro, Viale Europa, Catanzaro - 88100, Italy.
E-mail: gallelli@unicz.it

formulation of atenolol (Tenormin® 50 mg bid) to the generic one (atenolol Almus generics® 50 mg bid). About 14 days later the patient developed palpitations, and in November 2013, she was admitted to the Medicine Interne Operative Unit, with a diagnosis of atrial fibrillation. Both blood and urinary chemical analyses, to evaluate the activity of the kidney, liver, and thyroid, were negative. Similarly, the urinary catecholamine levels were normal. The drug levels were checked and it was demonstrated that her plasma levels of atenolol (45 ng/mL; normal range 50–70 ng/mL) were subtherapeutic. Given this temporal relationship of changing the Tenormin®-based therapy to the generic type of atenolol therapy could have led to the subtherapeutic beta-blocker levels, resulting in recurrence of atrial fibrillation, although this may be a causal relationship. The Naranjo probability scale^[4] documented a possible association between generic atenolol and heart symptoms, therefore, generic atenolol was switched to the brand formulation (Tenormin® 50 mg bid), with an initial improvement of symptoms in about three days and with complete control of symptoms in about 14 days (atenolol plasma levels: 57 ng/mL). The patient was also treated with dabigatran (Pradaxa®, 150 mg bid). During this time, the dosages of other medications, such as, tapazole and telmisartan, remained unchanged.

DISCUSSION

Beta blockers represent the initial treatment for rate control as well as maintenance of sinus rhythm in patients with paroxysmal atrial fibrillation.^[5] We have reported a case of recurrence of atrial fibrillation after a switch from the chronic treatment with Tenormin® to generic atenolol.

Several factors may be involved in the genesis of atrial fibrillation,^[6-9] however, in our patient clinical manifestation and laboratory findings excluded a suggestion that atrial fibrillation was related to other cardiac or systemic diseases.

Previously, we described that dispensing error may be involved in the development of side effects or in the lack of efficacy.^[10,11] In our patient, we can exclude an error in the timing of drug administration or in the drug used, as referred by the patient and his parents.

As it has been reported that drug–drug interaction may be able to induce clinical ineffectiveness,^[12] pharmacological evaluation has excluded that these could have played a role in the development of atrial fibrillation in our patient.

Using the Naranjo probability scale, we documented an association between the generic drug and the development of atrial fibrillation, and we demonstrated in this case that a switch from the generic formulation of atenolol to the Tenormin®-based therapy resulted in the improvement of

Table 1: Differences in excipients between the brand (Tenormin®) and generic (Atenolol Almus®) formulations of atenolol

Tenormin®	Atenolol Almus®
Heavy magnesium carbonate	Heavy magnesium carbonate
Corn starch	Corn starch
Sodium lauryl sulfate	Sodium lauryl sulfate
Gelatin	Magnesium stearate
Magnesium stearate	

symptoms, with an increase in plasma atenolol values. As other articles have suggested a difference in the clinical efficacy of the generic drug with respect to the brand formulations,^[13-16] in recent times, a retrospective study performed in Italy has reported that off-patent generic drugs have the same efficacy as the brand formulation.^[17]

Some factors such as differences in bioequivalence, excipients, and impurity may be involved in the difference of clinical efficacy between brand and generic formulations.^[15,16,18,19] However, in the present case we can exclude the role of excipients, because the generic formulation shows excipients similar to those in the brand formulation [Table 1].

In conclusion, as this is only a case report, we suggest that treatment must not be changed from a brand formulation to a generic one when the patient is chronically treated, or if it is necessary we suggest performing a plasma pharmacological evaluation of the drug concentration in order to avoid possible clinical ineffectiveness.

REFERENCES

- Gowd BM, Thompson PD. Effect of female sex on cardiac arrhythmias. *Cardiol Rev* 2012;20:297-303.
- Boateng S. Tachycardia. *Dis Mon* 2013;59:74-82.
- Naik A. Beta blockers in arrhythmias: When and where to use? *Indian Heart J* 2010;62:136-8.
- Naranjo CA, Busto U, Sellers EM, Sandor P, Ruiz I, Roberts EA, *et al.* A method for estimating the probability of adverse drug reactions. *Clin Pharmacol Ther* 1981;30:239-45.
- Gillis AM, Verma A, Talajic M, Nattel S, Dorian P; CCS Atrial Fibrillation Guidelines Committee. Canadian Cardiovascular Society atrial fibrillation guidelines 2010: Rate and rhythm management. *Can J Cardiol* 2011;27:47-59.
- Menezes AR, Lavie CJ, Dinicolantonio JJ, O'Keefe J, Morin DP, Khatib S, *et al.* Cardiometabolic risk factors and atrial fibrillation. *Rev Cardiovasc Med* 2013;14:e73-81.
- Menezes AR, Lavie CJ, DiNicolantonio JJ, O'Keefe J, Morin DP, Khatib S, *et al.* Atrial fibrillation in the 21st century: A current understanding of risk factors and primary prevention strategies. *Mayo Clin Proc* 2013;88:394-409.
- Balta S, Kurtoglu E, Demir M, Demirkol S, Arslan Z, Unlu M. Risk factors for new-onset atrial fibrillation. *Int J Cardiol* 2014;171:e46.
- Knuiman M, Briffa T, Divitini M, Chew D, Eikelboom J, McQuillan B, *et al.* A cohort study examination of established and emerging risk factors for atrial fibrillation: The Busselton Health Study. *Eur J Epidemiol* 2014;29:181-90.
- Di Mizio G, Gallelli L, Barbieri V, Loiacono D, Colosimo F, Rende P,

Gallelli, *et al.*: Generic atenolol and clinical ineffectiveness

- et al.* Capecitabine-induced, rapid decrease of renal function due to drug dispensing error in a hospital pharmacy. *J Clin Pharmacol* 2011;51:117-9.
11. Gallelli L, Staltari O, Palleria C, Di Mizio G, De Sarro G, Caroleo B. A case of adverse drug reaction induced by dispensing error. *J Forensic Leg Med* 2012;19:497-8.
 12. Palleria C, Di Paolo A, Giofrè C, Caglioti C, Leuzzi G, Siniscalchi A, *et al.* Pharmacokinetic drug-drug interaction and their implication in clinical management. *J Res Med Sci* 2013;18:601-10.
 13. Reiffel JA. Formulation substitution and other pharmacokinetic variability: Underappreciated variables affecting antiarrhythmic efficacy and safety in clinical practice. *Am J Cardiol* 2000;85 (10A):46-52D.
 14. Reiffel JA, Kowey PR. Generic antiarrhythmics are not therapeutically equivalent for the treatment of tachyarrhythmias. *Am J Cardiol* 2000;85:1151-3, A10.
 15. Gallelli L, Palleria C, De Vuono A, Mumoli L, Vasapollo P, Piro B, *et al.* Safety and efficacy of generic drugs with respect to brand formulation. *J Pharmacol Pharmacother* 2013;4(Suppl 1):S110-4.
 16. De Vuono A, Palleria C, Scicchitano F, Squillace A, De Sarro G, Gallelli L. Skin rash during treatment with generic itraconazole. *J Pharmacol Pharmacother* 2014;5:158-60.
 17. Colombo GL, Agabiti-Rosei E, Margonato A, Mencacci C, Montecucco CM, Trevisan R. Off-patent generic medicines vs. off-patent brand medicines for six reference drugs: A retrospective claims data study from five local healthcare units in the Lombardy Region of Italy. *PloS One* 2013;8:e82990.
 18. De Vuono A, Scicchitano F, Palleria C, Russo E, De Sarro G, Gallelli L. Lack of efficacy during the switch from brand to generic allopurinol. *J Forensic Leg Med* 2013;20:540-2.
 19. Yamamoto Y, Fukami T, Koide T, Onuki Y, Suzuki T, Metori K, *et al.* Comparative pharmaceutical evaluation of brand and generic clobetasone butyrate ointments. *Int J Pharm* 2014;463:62-7.

How to cite this article: Gallelli L, Maida F, Staltari O, Rende P, Russo E, Caroleo B, *et al.* Recurrence of atrial fibrillation after switching from brand to generic atenolol. *J Pharmacol Pharmacother* 2015;6:39-41.

Source of Support: Nil, **Conflict of Interest:** None declared.

Author Help: Online submission of the manuscripts

Articles can be submitted online from <http://www.journalonweb.com>. For online submission, the articles should be prepared in two files (first page file and article file). Images should be submitted separately.

1) First Page File:

Prepare the title page, covering letter, acknowledgement etc. using a word processor program. All information related to your identity should be included here. Use text/rtf/doc/pdf files. Do not zip the files.

2) Article File:

The main text of the article, beginning with the Abstract to References (including tables) should be in this file. Do not include any information (such as acknowledgement, your names in page headers etc.) in this file. Use text/rtf/doc/pdf files. Do not zip the files. Limit the file size to 1 MB. Do not incorporate images in the file. If file size is large, graphs can be submitted separately as images, without their being incorporated in the article file. This will reduce the size of the file.

3) Images:

Submit good quality color images. Each image should be less than 4096 kb (4 MB) in size. The size of the image can be reduced by decreasing the actual height and width of the images (keep up to about 6 inches and up to about 1800 x 1200 pixels). JPEG is the most suitable file format. The image quality should be good enough to judge the scientific value of the image. For the purpose of printing, always retain a good quality, high resolution image. This high resolution image should be sent to the editorial office at the time of sending a revised article.

4) Legends:

Legends for the figures/images should be included at the end of the article file.