

Case Report

Terlipressin-induced Peripheral Ischemic Gangrene in a Diabetic Patient

Phulen Sarma, Gaurav Muktesh¹, Narender Dhaka¹, Rakesh Ruhela, Abhishek Mishra, Rahul Singh¹, Saroj K. Sinha¹, Bikash Medhi, Rakesh Kochhar¹

Departments of Pharmacology and ¹Gastroenterology, Postgraduate Institute of Medical Education and Research, Chandigarh, India

Abstract

Terlipressin is used commonly in the management of hepatorenal syndrome and acute variceal bleeding. Like its parent compound vasopressin, it is also notorious for its ischemic complications. Terlipressin-induced ischemic complications can virtually affect any part of the body, but the incidence of serious complications is less than its parent compound vasopressin. Here, we report a case of terlipressin-induced peripheral ischemic gangrene in a diabetic male, which ultimately led to death of the patient.

Keywords: Gangrene, hepatorenal syndrome, ischemic complication, terlipressin, variceal bleeding

INTRODUCTION

Acute variceal bleed and hepatorenal syndrome are the two important causes of morbidity and mortality in patients with cirrhosis. Acute bleeding esophageal varices is associated with 10%–50% mortality per episode.^[1,2] Treatment options are pharmacotherapy (splanchnic vasoconstrictors, antibiotics), endoscopic therapy (ligation, sclerotherapy), other salvage procedures (balloon tamponade, transjugular intrahepatic portosystemic shunt, and balloon occluded transvenous retrograde obliteration), and surgical therapies such as distal splenorenal shunt, angiectomy, and liver transplantation. Among pharmacotherapeutic options are vasopressin and its analog (terlipressin) and somatostatin and its analogs (vapeotide and octreotide).^[3] Pharmacotherapeutic interventions need comparatively less expertise than endoscopic procedures, but controversy exists regarding their risk–benefit ratio.^[1] Vasopressin is the most potent splanchnic vasoconstrictor, but increased incidence of ischemic side effects limits its use. Terlipressin is a vasopressin analog with longer duration of action with fewer ischemic complications.^[4] Vasoactive therapy is associated with significantly lower risk of acute all-cause mortality, transfusion requirements, improved control of bleeding and shorter hospital stay.^[1] Although side effect profile of terlipressin is more favorable, compared to parent compound vasopressin, yet few cases are reported regarding ischemic complications of terlipressin therapy.^[5–11]

Ischemia leading to gangrene is a relatively rare adverse effect of terlipressin. Only few cases are reported till now. Here, we report a case of terlipressin-induced peripheral gangrene in a 65-year-old male patient with acute variceal bleed.

CASE REPORT

A 65-year-old male diabetic (fasting blood sugar 166 mg/dl, postprandial blood sugar 240 mg/dl, HbA1c 8.1%) patient presented in emergency with complaints of hematemesis and melena for the past 24 h. He had been a chronic alcoholic consuming around 100 g of alcohol per day for the last 35 years. On evaluation, he was found to have hepatomegaly. His hemogram showed anemia (8.9 g/dl), thrombocytopenia (platelet count 0.4 lac/mm³) with evidence of coagulopathy (prothrombin time index 48%). Ultrasound abdomen showed altered echotexture of liver with features of portal hypertension. On the basis of clinical, biochemical, and radiological imaging, provisional diagnosis of chronic liver disease with upper gastrointestinal (GI) bleed (variceal) was made and managed with blood transfusion, antibiotics,

Address for correspondence: Bikash Medhi,
Department of Pharmacology, Postgraduate Institute of Medical Education
and Research, Chandigarh - 160 012, India.
E-mail: drbikashus@yahoo.com

Access this article online

Quick Response Code:



Website:
www.jpharmacol.com

DOI:
10.4103/jpp.JPP_42_17

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Sarma P, Muktesh G, Dhaka N, Ruhela R, Mishra A, Singh R, *et al.* Terlipressin-induced peripheral ischemic gangrene in a diabetic patient. *J Pharmacol Pharmacother* 2017;8:148-50.

Received: 18-03-2017 **Revised:** 16-04-2017 **Accepted:** 09-08-2017

and splanchnic vasoconstrictor (injection terlipressin 1 mg 6 hourly). After initial stabilization, the patient underwent upper GI endoscopy which showed large esophageal varices, for which endoscopic variceal ligation was done. After 2 days of therapy, he developed blackening of bilateral upper and lower limbs (UL and LL) which was more prominent in distal parts [Figures 1-3]. This blackening evolved rapidly into large necrotizing areas with fluid exudation. On examination, his distal arterial pulses were palpable. Color Doppler study showed normal color flow and spectral waveform in all major arteries of bilateral UL and LL. X-ray of both bilateral UL and LL showed normal study.

Immediately, terlipressin was thought to be the culprit drug for this complication and was discontinued. After that, the proximal progression of discoloration stopped. Meanwhile, the patient was screened for markers of autoimmunity in the form of antinuclear antibody and antineutrophil cytoplasm antibody which were found to be negative.

He was given symptomatic treatment in the form of vigorous dressing, antibiotics, and intravenous fluids. He was offered amputation of bilateral toes and fingers to prevent further progression of the disease and resultant sepsis. However, the patient refused for an amputation and he went against medical advice. After 15 days, he died of sepsis at his home.

On causality assessment, by both WHO-UMC^[12] and Naranjo scale,^[13] terlipressin was found to be of probable category by both the classification systems while other concomitant medications were found to be of unlikely causality. Severity was assessed by modified Hartwig and Siegel Scale,^[14] and it was found to be of severe category. Seriousness was assessed by US-Food and Drug Administration criteria^[15] and it was a serious event. Preventability of the adverse drug reaction (ADR) was assessed by Schumock and Thronton's scale.^[16] Moreover, this ADR was found to be of definitely not preventable category. Predictability analysis was done by predictability scale by Imbs *et al.*,^[17] and it was found to be of definitely not predictable (Category C).

DISCUSSION

Ischemic complications are relatively rare adverse effects of terlipressin therapy. In our case, the patient developed peripheral gangrene and succumbed to it. The spectrum of ischemic complication ranges from peripheral gangrene, myocardial ischemia, and ischemia of bowel, and the spectrum of ischemic complications practically involved every organ extending from scalp ischemia to ischemic complications of the scrotum.^[5-11,18-21] Most cases were managed conservatively with stopping of terlipressin.^[5-11,18-21] In two cases, sildenafil was tried as an intervention and one case recorded significant improvement and recovered completely at 1 month, while other case developed decreased urine output and alprostadil and sildenafil had to be withdrawn. Although amputation had to be done on medical grounds, this patient also recovered.^[5,6] V1 receptor-mediated vasoconstriction and superimposed



Figure 1: After 2 days of therapy, blackening of the upper limb



Figure 2: After 2 days of therapy, blackening of the lower limb



Figure 3: After 2 days of therapy, blackening of the lower and upper limbs

V2 receptor-mediated Von Will brand factor release from endothelial cells, together are implicated in the pathogenesis of ischemic complications.^[22-24] Role of pharmacogenomics in vasopressin-induced ischemic complications was evaluated by Anantasit *et al.* and a significant association was found between serious adverse effects of vasopressin and

single-nucleotide polymorphism of rs28418396, implicating a role of pharmacogenomics in the prediction of ischemic complications.^[23] This area needs further study. We report this rare adverse effect of terlipressin so that physicians will be aware of these dreaded adverse effects of terlipressin as timely detection and stopping of terlipressin at the earliest sign of ischemia can be lifesaving.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Wells M, Chande N, Adams P, Beaton M, Levstik M, Boyce E, *et al.* Meta-analysis: Vasoactive medications for the management of acute variceal bleeds. *Aliment Pharmacol Ther* 2012;35:1267-78.
- Dobre M, Demirjian S, Sehgal AR, Navaneethan SD. Terlipressin in hepatorenal syndrome: A systematic review and meta-analysis. *Int Urol Nephrol* 2011;43:175-84.
- Kim YD. Management of acute variceal bleeding. *Clin Endosc* 2014;47:308-14.
- Garcia-Tsao G, Sanyal AJ, Grace ND, Carey WD, Practice Guidelines Committee of American Association for Study of Liver Diseases. Practice Parameters Committee of American College of Gastroenterology. *et al.* Prevention and management of gastroesophageal varices and variceal hemorrhage in cirrhosis. *Am J Gastroenterol* 2007;102:2086-102.
- Lee HJ, Oh MJ. A case of peripheral gangrene and osteomyelitis secondary to terlipressin therapy in advanced liver disease. *Clin Mol Hepatol* 2013;19:179-84.
- Bañuelos Ramírez DD, Sánchez Alonso S, Ramírez Palma MM. Sildenafil in severe peripheral ischemia induced by terlipressin. A case report. *Reumatol Clin* 2011;7:59-60.
- Lee MY, Chu CS, Lee KT, Lee HC, Su HM, Cheng KH, *et al.* Terlipressin-related acute myocardial infarction: A case report and literature review. *Kaohsiung J Med Sci* 2004;20:604-8.
- Ghatak T, Poddar B, Mahindra S. Left ventricular failure and left ventricular inferior wall hypokinesia following terlipressin injection. *Ann Card Anaesth* 2014;17:257-9.
- Mégarbané H, Barete S, Khosrotehrani K, Izzedine H, Moguelet P, Chosidow O, *et al.* Two observations raising questions about risk factors of cutaneous necrosis induced by terlipressin (Glypressin). *Dermatology* 2009;218:334-7.
- Kim HR, Lee YS, Yim HJ, Lee HJ, Ryu JY, Lee HJ, *et al.* Severe ischemic bowel necrosis caused by terlipressin during treatment of hepatorenal syndrome. *Clin Mol Hepatol* 2013;19:417-20.
- Willems MG, Schoenemann J, Rey C, Schäfer H, Lindeken KD. Ischemia of the cecum caused by glycyllipressin. *Leber Magen Darm* 1985;15:165-8.
- The Use of the WHO-UMC System for Standardized Case Causality Assessment: The Uppsala Monitoring Center. Available from: <http://www.who-umc.org/Graphics/24734.pdf>. [Last accessed on 2016 Feb 22].
- Naranjo CA, Busto U, Sellers EM, Sandor P, Ruiz I, Roberts EA, *et al.* A method for estimating the probability of adverse drug reactions. *Clin Pharmacol Ther* 1981;30:239-45.
- Hartwig SC, Siegel J, Schneider PJ. Preventability and severity assessment in reporting adverse drug reactions. *Am J Hosp Pharm* 1992;49:2229-32.
- What is a Serious Adverse Event? Available from: <http://www.fda.gov/Safety/MedWatch/HowToReport/ucm053087.htm>. [Last accessed on 2016 Feb 22].
- Schumock GT, Thornton JP. Focusing on the preventability of adverse drug reactions. *Hosp Pharm* 1992;27:538.
- Imbs JL, Pletan Y, Spriet A. Assessment of preventable iatrogenic drug therapy: Methodology. Round Table No 2 at Giens XIII. *Therapie* 1998;53:365-70.
- Lee JS, Lee HS, Jung SW, Han WS, Kim MJ, Lee SW, *et al.* A case of peripheral ischemic complication after terlipressin therapy. *Korean J Gastroenterol* 2006;47:454-7.
- Sundriyal D, Kumar N, Patnaik I, Kamble U. Terlipressin induced ischaemia of skin. *BMJ Case Rep* 2013;2013. pii: ber2013010050.
- Chandail VS, Jamwal V. Cutaneous complications of terlipressin. *JK Sci J Med Educ Res* 2011;13:205-7.
- Posada C, Feal C, García-Cruz A, Alvarez V, Alvarez M, Cruces MJ, *et al.* Cutaneous necrosis secondary to terlipressin therapy. *Acta Derm Venereol* 2009;89:434-5.
- Kam PC, Williams S, Yoong FF. Vasopressin and terlipressin: Pharmacology and its clinical relevance. *Anaesthesia* 2004;59:993-1001.
- Anantasi N, Boyd JH, Walley KR, Russell JA. Serious adverse events associated with vasopressin and norepinephrine infusion in septic shock. *Crit Care Med* 2014;42:1812-20.
- Chang YH, Yen FC, Hsieh MC, Lin KD, Shin SJ, Hsin SC, *et al.* Diabetic muscle infarction in association with terlipressin therapy: A case report. *Kaohsiung J Med Sci* 2009;25:25-8.